
PLANT STANOLS PROVIDE ADDITIONAL CHOLESTEROL-LOWERING ALONGSIDE STATINS




European guidelines recommend plant

stanols as part of the management of raised cholesterol as an adjunct to drug therapy in high and very high risk patients who fail to achieve LDL cholesterol target levels on statins or are statin intolerant



The distinct mechanism of action of plant stanols for **reducing cholesterol absorption in the intestine** allows an additive effect to statins



...discontinue statins within 1 year of treatment initiation, and more do so over longer time periods⁸ Adherence to statin treatment has been reported to be poor in both the short and long term, emphasising the importance of maintaining cholesterol lowering dietary measures alongside medication⁹

Plant stanol ester has been shown to lower cholesterol. High cholesterol is a risk factor in the development of coronary heart disease. A daily intake of 1.5-2.4 g plant stanols lower cholesterol by 7-10% in 2 to 3 weeks. The beneficial effect is obtained with a daily intake of 1.5 to 3 g plant stanols.

LDL: low-density lipoprotein.

References:


World Health Organization. Raised cholesterol. www.who.int/gho/ncd/risk_factors/cholesterol_text/en/
Scholle JM, et al. J Am Coll Nutr. 2009; 28(5): 517-524. 3. Gylling H, et al. Atherosclerosis 2014; 232(2): 346-360. 4. Hallikainen M, et al. Atherosclerosis 2011; 217(2): 473-478. 5. Blair SN, et al. Am J Cardiol 2000; 86(1): 6-52. 6. de Jong A, et al. Br J Nutr 2008; 100(5): 937-941. 7. Mach F, Baigent C, Catapano AL, et al. Eur Heart J 2019; doi:10.1093/eurheartj/ehz455. 8. Mikhailidis DP, et al. Curr Med Res Opin 2016; 32(10): 1639-1640. 9. Ofori-Asenso R, et al. J Gerontol A Biol Sci Med Sci 2018; 73(6): 813-819.



PLANT STANOL ESTER

Fast and sustainable cholesterol reduction

Maintaining a healthy blood cholesterol level is important to support heart wellbeing. Plant stanol ester, the cholesterol-lowering ingredient in **Benecol® functional foods**, provides an effective way to lower LDL cholesterol as part of a healthy diet



Benecol offers a broad range of products to ensure that achieving adequate plant stanol intake is easy



*Plant stanol ester has been shown to lower cholesterol. High cholesterol is a risk factor in the development of coronary heart disease. A daily intake of 1.5–2.4 g plant stanols lowers cholesterol by 7–10% in 2 to 3 weeks.

Please visit www.benecol.co.uk/hcp (UK) or www.benecol.ie/hcp (IE) for more information on plant stanol ester for healthcare professionals.

For more information on our clinical studies or for patient leaflets and coupons please call 0800 018 4010 (UK) 1800 551 707 (ROI) (Mon-Fri 09:00-17:00)

LDL: low-density lipoprotein. References:

Mach F, Baigent C, Catapano AL *et al.* Eur Heart J 2019; doi:10.1093/eurheartj/ehz455.
Piepoli MF, Hoes AW, Agewall S *et al.* Eur Heart J 2016; 37: 2315–2381.

